

d his

LYCOOK
11/22/04

(FILE 'HOME' ENTERED AT 22:30:07 ON 22 NOV 2004)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
22:30:28 ON 22 NOV 2004

L1	0 S (CB 10 PEPTIDE)
L2	216 S (CB10)
L3	115 S L2 AND PEPTIDE?
L4	100 S L3 AND COLLAGEN?
L5	34 S L4 AND ANTIB?
L6	2 S L5 AND KIT?
L7	2 DUPLICATE REMOVE L6 (0 DUPLICATES REMOVED)
L8	16 DUPLICATE REMOVE L5 (18 DUPLICATES REMOVED)

=>

on STN

AN 91106346 EMBASE

DN 1991106346

TI The role of **collagen** conformation in type II anticollagen immunity in rheumatoid arthritis.

AU Boissier M.C.; Chiocchia G.; Fournier C.

CS Departement de Rhumatologie, Hopital Avicenne, Bobigny, France

SO Revue du Rhumatisme et des Maladies Osteo-Articulaires, (1991) 58/1 (19-24).

ISSN: 0035-2659 CODEN: RRMOA2

CY France

DT Journal; Article

FS 031 Arthritis and Rheumatism

LA French

SL English; Spanish; German

AB Type II anticollagen (CII) autoimmunity is a frequently reported, but non-specific, phenomenon in rheumatoid arthritis (RA). The authors show that in 88 sera samples from patients suffering from RA, the incidence of **antibodies** targeted against endogenous human CII was the same as that found for 149 control blood donors (14.8% versus 11.4%). However, a significant difference was found for the incidence of **antibodies** targeted against the α -chains of CII (26.1% versus 6.0%, $p < 0.001$). As a result of investigating the specificity of the anti-CII **antibodies** in greater detail by means of an immunoprinting of the CII **peptide** fragments obtained after splitting the molecule by cyanogen bromide, the authors have demonstrated that the largest CII **peptides** (CB10 and CB11) were better recognized than the smaller **peptides** (CB8, CB9.7), with no significant difference between PR and control plasmas. Using competitive methods, evidence was obtained in support of heterogeneous recognition by the anti-CII **antibodies**: some recognize conformational determinants only, whereas others are targeted against the primary sequences of the α -1 (II) chain.

CT Medical Descriptors:

*rheumatoid arthritis: ET, etiology

adult

article

autoimmunity

controlled study

female

human

human tissue

immunoblotting

major clinical study

male

Drug Descriptors:

collagen type 2

***antibody**

AN 91106346 EMBASE

DN 1991106346

TI The role of **collagen** conformation in type II anticollagen immunity in rheumatoid arthritis.

AU Boissier M.C.; Chiocchia G.; Fournier C.

CS Departement de Rhumatologie, Hopital Avicenne, Bobigny, France

SO Revue du Rhumatisme et des Maladies Osteo-Articulaires, (1991) 58/1 (19-24).

ISSN: 0035-2659 CODEN: RRM0A2

CY France

DT Journal; Article

FS 031 Arthritis and Rheumatism

LA French

SL English; Spanish; German

AB Type II anticollagen (CII) autoimmunity is a frequently reported, but non-specific, phenomenon in rheumatoid arthritis (RA). The authors show that in 88 sera samples from patients suffering from RA, the incidence of **antibodies** targeted against endogenous human CII was the same as that found for 149 control blood donors (14.8% versus 11.4%). However, a significant difference was found for the incidence of **antibodies** targeted against the α -chains of CII (26.1% versus 6.0%, $p < 0.001$). As a result of investigating the specificity of the anti-CII **antibodies** in greater detail by means of an immunoprinting of the CII **peptide** fragments obtained after splitting the molecule by cyanogen bromide, the authors have demonstrated that the largest CII **peptides** (CB10 and CB11) were better recognized than the smaller **peptides** (CB8, CB9.7), with no significant difference between PR and control plasmas. Using competitive methods, evidence was obtained in support of heterogeneous recognition by the anti-CII **antibodies**: some recognize conformational determinants only, whereas others are targeted against the primary sequences of the α -1 (II) chain.

CT Medical Descriptors:

*rheumatoid arthritis: ET, etiology

adult

article

autoimmunity

controlled study

female

human

human tissue

immunoblotting

major clinical study

male

Drug Descriptors:

collagen type 2

***antibody**

AN 1992:455101 BIOSIS

DN PREV199294096501; BA94:96501

TI **COLLAGEN**-INDUCED ARTHRITIS IN RATS EXAMINATION OF THE EPITOPE
SPECIFICITIES OF CIRCULATING AND CARTILAGE-BOUND **ANTIBODIES**
PRODUCED BY OUTBRED AND INBRED RATS USING CYANOGEN BROMIDE-DERIVED
PEPTIDES PURIFIED FROM HETEROLOGOUS AND HOMOLOGOUS TYPE II
COLLAGENS.

AU CREMER M A [Reprint author]; TERATO K; WATSON W C; GRIFFITHS M M; TOWNES A
S; KANG A H

CS RES SERV 151 , VETERANS ADM MED CENT, 1030 JEFFERSON AVE, MEMPHIS, TENN
38104, USA

SO Journal of Immunology, (1992) Vol. 149, No. 3, pp. 1045-1053.
CODEN: JOIMA3. ISSN: 0022-1767.

DT Article

FS BA

LA ENGLISH

ED Entered STN: 7 Oct 1992

Last Updated on STN: 8 Oct 1992

AB To determine the number and location of **antibody** binding
epitopes on type II **collagen**, outbred and inbred rats were
immunized with chick, bovine, human, and rat type II **collagen**
(CII, BII, HII, and RII); all sera were assayed for reaction with a panel
of CB **peptides** purified and renatured from the immunizing
collagen and from RI. **Antibody** reaction patterns
(profiles) varied among individual outbred rats but were essentially
constant over time and changed little after boosting. The strongest
antibody reactions were to CB11, CB9-7, and CB12 followed by CB8,
CB10, and CB6. **Antibody** profiles varied depending on
the species of **collagen** used for immunization and the strain of
rat immunized. Except for CB10, where **antibodies** were
largely specific for heterologous **collagens**, **antibodies**
reactive with all other CB **peptides** cross-reacted strongly with
renatured rat CB **peptides**. Sera from inbred BB rats immunized
with BII, CII, or HII reacted best with CB11, unlike antisera to RII that
reacted strongly with CB9-7. Inbred LEW, COP, WKY, F344, and BUF rats
immunized with BII reacted strongest with CB9-7 and variably with CB11 and
CB 12. BBxLEW F1 hybrid rats reacted almost equally with CB11 and CB9-7
producing an **antibody** profile intermediate to those elicited in
the parent strains. Finally, **antibodies** reactive with rat CB11,
CB9-7, and CB12 could be eluted from normal rat cartilage incubated in
anti-BII serum; **antibody** eluate profiles generally paralleled
the profile produced by the sera applied to cartilage. Taken together,
these findings indicate that multiple **antibody**-reactive epitopes
on type II **collagen** may be instrumental in the initiation of
collagen-induced arthritis in rats, particularly shared or
cross-reactive epitopes located within CB11, CB9-7, CB12, and CB6.

CC Genetics - Animal 03506

Biochemistry studies - Proteins, peptides and amino acids 10064

Pathology - Inflammation and inflammatory disease 12508

Bones, joints, fasciae, connective and adipose tissue - Pathology 18006

Immunology - Immunopathology, tissue immunology 34508

IT Major Concepts

Genetics; Immune System (Chemical Coordination and Homeostasis);

Skeletal System (Movement and Support)

ORGN Classifier

Muridae 86375

Super Taxa

Rodentia; Mammalia; Vertebrata; Chordata; Animalia

Taxa Notes

Animals, Chordates, Mammals, Nonhuman Vertebrates, Nonhuman Mammals,

AN 1992:455101 BIOSIS
DN PREV199294096501; BA94:96501
TI **COLLAGEN**-INDUCED ARTHRITIS IN RATS EXAMINATION OF THE EPITOPE
SPECIFICITIES OF CIRCULATING AND CARTILAGE-BOUND **ANTIBODIES**
PRODUCED BY OUTBRED AND INBRED RATS USING CYANOGEN BROMIDE-DERIVED
PEPTIDES PURIFIED FROM HETEROLOGOUS AND HOMOLOGOUS TYPE II
COLLAGENS.
AU CREMER M A [Reprint author]; TERATO K; WATSON W C; GRIFFITHS M M; TOWNES A
S; KANG A H
CS RES SERV 151 , VETERANS ADM MED CENT, 1030 JEFFERSON AVE, MEMPHIS, TENN
38104, USA
SO Journal of Immunology, (1992) Vol. 149, No. 3, pp. 1045-1053.
CODEN: JOIMA3. ISSN: 0022-1767.
DT Article
FS BA
LA ENGLISH
ED Entered STN: 7 Oct 1992
Last Updated on STN: 8 Oct 1992
AB To determine the number and location of **antibody** binding
epitopes on type II **collagen**, outbred and inbred rats were
immunized with chick, bovine, human, and rat type II **collagen**
(CII, BII, HII, and RII); all sera were assayed for reaction with a panel
of CB **peptides** purified and renatured from the immunizing
collagen and from RI. **Antibody** reaction patterns
(profiles) varied among individual outbred rats but were essentially
constant over time and changed little after boosting. The strongest
antibody reactions were to CB11, CB9-7, and CB12 followed by CB8,
CB10, and CB6. **Antibody** profiles varied depending on
the species of **collagen** used for immunization and the strain of
rat immunized. Except for CB10, where **antibodies** were
largely specific for heterologous **collagens**, **antibodies**
reactive with all other CB **peptides** cross-reacted strongly with
renatured rat CB **peptides**. Sera from inbred BB rats immunized
with BII, CII, or HII reacted best with CB11, unlike antisera to RII that
reacted strongly with CB9-7. Inbred LEW, COP, WKY, F344, and BUF rats
immunized with BII reacted strongest with CB9-7 and variably with CB11 and
CB 12. BBxLEW F1 hybrid rats reacted almost equally with CB11 and CB9-7
producing an **antibody** profile intermediate to those elicited in
the parent strains. Finally, **antibodies** reactive with rat CB11,
CB9-7, and CB12 could be eluted from normal rat cartilage incubated in
anti-BII serum; **antibody** eluate profiles generally paralleled
the profile produced by the sera applied to cartilage. Taken together,
these findings indicate that multiple **antibody**-reactive epitopes
on type II **collagen** may be instrumental in the initiation of
collagen-induced arthritis in rats, particularly shared or
cross-reactive epitopes located within CB11, CB9-7, CB12, and CB6.
CC Genetics - Animal 03506
Biochemistry studies - Proteins, peptides and amino acids 10064
Pathology - Inflammation and inflammatory disease 12508
Bones, joints, fasciae, connective and adipose tissue - Pathology 18006
Immunology - Immunopathology, tissue immunology 34508
IT Major Concepts
Genetics; Immune System (Chemical Coordination and Homeostasis);
Skeletal System (Movement and Support)
ORGN Classifier
Muridae 86375
Super Taxa
Rodentia; Mammalia; Vertebrata; Chordata; Animalia
Taxa Notes
Animals, Chordates, Mammals, Nonhuman Vertebrates, Nonhuman Mammals,

AN 1992:455101 BIOSIS

DN PREV199294096501; BA94:96501

TI **COLLAGEN**-INDUCED ARTHRITIS IN RATS EXAMINATION OF THE EPITOPE
SPECIFICITIES OF CIRCULATING AND CARTILAGE-BOUND **ANTIBODIES**
PRODUCED BY OUTBRED AND INBRED RATS USING CYANOGEN BROMIDE-DERIVED
PEPTIDES PURIFIED FROM HETEROLOGOUS AND HOMOLOGOUS TYPE II
COLLAGENS.

AU CREMER M A [Reprint author]; TERATO K; WATSON W C; GRIFFITHS M M; TOWNES A
S; KANG A H

CS RES SERV 151 , VETERANS ADM MED CENT, 1030 JEFFERSON AVE, MEMPHIS, TENN
38104, USA

SO Journal of Immunology, (1992) Vol. 149, No. 3, pp. 1045-1053.
CODEN: JOIMA3. ISSN: 0022-1767.

DT Article

FS BA

LA ENGLISH

ED Entered STN: 7 Oct 1992

Last Updated on STN: 8 Oct 1992

AB To determine the number and location of **antibody** binding
epitopes on type II **collagen**, outbred and inbred rats were
immunized with chick, bovine, human, and rat type II **collagen**
(CII, BII, HII, and RII); all sera were assayed for reaction with a panel
of CB **peptides** purified and renatured from the immunizing
collagen and from RI. **Antibody** reaction patterns
(profiles) varied among individual outbred rats but were essentially
constant over time and changed little after boosting. The strongest
antibody reactions were to CB11, CB9-7, and CB12 followed by CB8,
CB10, and CB6. **Antibody** profiles varied depending on
the species of **collagen** used for immunization and the strain of
rat immunized. Except for CB10, where **antibodies** were
largely specific for heterologous **collagens**, **antibodies**
reactive with all other CB **peptides** cross-reacted strongly with
renatured rat CB **peptides**. Sera from inbred BB rats immunized
with BII, CII, or HII reacted best with CB11, unlike antisera to RII that
reacted strongly with CB9-7. Inbred LEW, COP, WKY, F344, and BUF rats
immunized with BII reacted strongest with CB9-7 and variably with CB11 and
CB 12. BBxLEW F1 hybrid rats reacted almost equally with CB11 and CB9-7
producing an **antibody** profile intermediate to those elicited in
the parent strains. Finally, **antibodies** reactive with rat CB11,
CB9-7, and CB12 could be eluted from normal rat cartilage incubated in
anti-BII serum; **antibody** eluate profiles generally paralleled
the profile produced by the sera applied to cartilage. Taken together,
these findings indicate that multiple **antibody**-reactive epitopes
on type II **collagen** may be instrumental in the initiation of
collagen-induced arthritis in rats, particularly shared or
cross-reactive epitopes located within CB11, CB9-7, CB12, and CB6.

CC Genetics - Animal 03506

Biochemistry studies - Proteins, peptides and amino acids 10064

Pathology - Inflammation and inflammatory disease 12508

Bones, joints, fasciae, connective and adipose tissue - Pathology 18006

Immunology - Immunopathology, tissue immunology 34508

IT Major Concepts

Genetics; Immune System (Chemical Coordination and Homeostasis);

Skeletal System (Movement and Support)

ORGN Classifier

Muridae 86375

Super Taxa

Rodentia; Mammalia; Vertebrata; Chordata; Animalia

Taxa Notes

Animals, Chordates, Mammals, Nonhuman Vertebrates, Nonhuman Mammals,